



*Opioid Agonist Therapy 101:
An Introduction to Clinical Practice Workshop*

Lindy's Tool

Marina Reinecke MBChB, CCFP(AM), ISAM

A dark blue arrow points to the right from the left edge of the slide. Below it, several thin, curved lines in shades of blue and grey sweep across the left side of the slide.

Faculty/Presenter Disclosure

- **Faculty:** Marina Reinecke
- **Relationships with commercial interests:** None



LEARNING OBJECTIVES

At the end of this educational activity, the learning will be able to:

- Discuss a practical tool for guiding a treatment conversation with an individual with opioid use disorder.
- List additional educational resources to further support decision making for your patient and his/her support people.



Choices in Opioid Addiction

1. Continue Using

2. Abstinence

3. Buprenorphine/Methadone

Addiction:

1. Loss of control or fighting for control.
2. Craving and compulsion to find and use your drug – no longer interested in previous activities and work.
3. Negative consequences, but can't stop using.

Choices in Opioid Addiction

1.

Continue Using

Most often,

↑ Use

↓ Control

↑ Consequences

Harm to Relationships

2.

Abstinence

Requires Time, Change, Practice

Requires:

- Knowledge of Self and Relationships
- Knowledge of Addiction
- New Habits and Coping Skill
- New Values and Perspective on Life
- May require Residential Treatment
- Taking care of yourself; service to others
- Practice and Learning
- Avoid Old/Slippery Friends, Places, Things, Habits, Alcohol, other Drugs

3.

Methadone/Buprenorphine

Stay on Opioid Agonist Therapy for Months or Years

Choices after Time

Stay on Methadone or Buprenorphine Indefinitely at same or lower dose

Taper off Slowly

Consider Abstinence When:


- Patient is young
- Oral opioids; no injection use
- Use < 6 -12 months
- Stable supportive family/work environment
- Patient's choice
- Patient's circumstances (finances, rural environment, etc.)
- < 30mgs of morphine equivalent per day

Consider buprenorphine/methadone when:

- Relapses after past treatment
- Injection use
- Polysubstance abuse
- Social stressors and instability
- Chronic pain complicated by addiction
- Patient's choice/patient circumstances
- Significant physical health problems from drug abuse
- Pregnancy
- Addicted family members
- >/= 30mgs morphine equivalent

RESOURCES

- ▶ 'Making the Choice, Making It Work; Treatment for Opioid Addiction' (CAMH)
- ▶ Buprenorphine/naloxone booklet – Chris A Cavacuiti, MD
- ▶ Buprenorphine/naloxone pamphlet
and
- ▶ Treatment program pamphlet from
Addictions Program, HSC



A dark grey arrow points right from the top left corner. Below it, several thin, curved lines in shades of blue and grey sweep across the left side of the slide. The background is a light blue gradient.

➔ Questions?