



*Opioid Agonist Therapy 101:  
An Introduction to Clinical Practice Workshop*

# **Withdrawal of Treatment**

**Marina Reinecke MBChB, CCFP(AM), ISAM**



# Faculty/Presenter Disclosure

- **Faculty:** Marina Reinecke
- **Relationships with commercial interests:** None



# LEARNING OBJECTIVES

At the end of this learning activity, the participant will be able to:

- Discuss a practical approach to the 3 most common scenarios involving withdrawal of treatment
- Discuss a practical approach to re-entry into treatment after involuntary withdrawal of treatment.



## 3 Scenarios

- 1. Voluntary – patient and clinic staff agree it is time to taper and attempt to stop treatment
- 2. Voluntary – Patient insists on tapering off although clinic staff are concerned
- 3. Involuntary – Staff decide that for reasons of safety (for the patient, public or staff) that the patient should not continue in treatment



# Scenario 1 – Ready to taper

- 1. social stability
- 2. emotional stability
- 3. no acute physical/psychiatric concerns
- 4. has left drug-involved life behind: no cocaine or methamphetamines, no benzodiazepines, no street access, support from non-using family and friends
- 5. recognizes risks and benefits of taper and plans accordingly
- 6. recovery supports – honesty, 12 steps, demonstrated stress tolerance strategies

# Scenario 1 – Ready to taper

- 1. discuss potential problems and how the taper might feel
- 2. together decide on pace of step-down; **small steps!!**
  - i) methadone 2 – 5 mg every few weeks
  - ii) buprenorphine/naloxone 2mg at a time ...or less!
  - iii) Sublocade – watch and wait; ? low dose  
buprenorphine/naloxone vs Probuphine implant

**Pay attention to the overall dose/picture**

- 3. patient and staff assess progress and watch for problems
- 4. Patient can phone to stop taper or to increase dose to re-stabilize

A decorative graphic on the left side of the slide. It features a dark blue vertical bar on the far left. A black arrow points to the right from the top of this bar. Several thin, light blue lines curve downwards and to the right from the base of the arrow, creating a sense of movement and flow.

# Denise

- As a teen, shy and lonely, eating disorder
- Was living with grandparents and grandpa was put on palliative care morphine – Denise started using his meds by injection
- Stabilized nicely at methadone clinic, really liked NA, did counselling



# Denise's taper

- Tapered down slowly from 90 to 35 mg. without complaint, looked well
- Did not discuss with clinic staff that she had started injecting intermittently
- Admitted to hospital with staph aureus empyema and septic hip
- Restabilized, treated this as a learning experience and tapering again





# Maureen

- Wild teenager with extreme polysubstance abuse, multiple overdoses, cutting – on Volume 4 of HSC chart
- Started to inject IV opioids
- Was clearly treated as an adult, with no positive response to borderline behaviour
- Did better than expected in treatment

A dark blue arrow points to the right from the left edge of the slide. Below it, several thin, curved lines in shades of blue and grey sweep across the left side of the slide.

## Maureen's taper

- After three years in program – had done lots of “growing up” and learned many new life skills
- Successfully tapered off over 14 months and still doing well

# Josh's taper

- ▶ Josh was a corrections officer who started using opioids for chronic MSK pain related to many sports injuries and tough workout routines like “punching a bag”.
- ▶ Stabilized on 24mgs of buprenorphine/naloxone and worked hard at dealing with the stress of work, an emotionally unstable X-wife, supporting his children financially and helping with maintenance projects around his parents' home (they took him in) over two years.
- ▶ Slowly tapered down by 2mgs increments every 1-2 months until he reached 10mgs; then 1mg increments down to 5mgs

## Josh's taper

- By now carrying a 1-month supply of buprenorphine/naloxone as a result of long-term stability.
- 3 incidents of “the cat knocking pill bottle into sink” in bathroom.
- 1<sup>st</sup> time – replaced; 2<sup>nd</sup> time replaced (witnessed) and reviewed safe storage etc. at in-person appointment. Denied increasing withdrawal or overuse.
- 3<sup>rd</sup> time replaced (witnessed in context of safety sensitive profession) and pulled carries. Looked well; no missed work etc.
- Watch and wait...



## End of Taper

- Addition is chronic illness – is patient prepared with plans to address emotional crisis, or if medical illness requires opioids?
- Consider offering follow-up for 3 – 6 months after methadone or buprenorphine/naloxone treatment ends.
- Review risk of cross - addiction (alcohol, cocaine etc..)



## Insists on Taper - ?Not Ready

- Slow taper is best but some patients will insist on rapid taper
- Discuss risks with patient repeatedly
- The patient will often refuse to recognize that problematic behaviour is occurring – e.g. - ongoing use of opioids or cocaine
- patient has right to decide



# Tapering, ? Not Ready

- Recognize when to stop the struggle
- Instead focus on your partnership
- Remind patient that it is never too late to “change your mind”
- There may be value to letting the patient “crash and burn” – patient hopefully returns to program with different expectations and behaviour
- Consider need for behaviour agreement if accepting back into treatment – raise the bar



## John's Taper

- He had failed 4 tries at abstinence and sadly started methadone – immediately felt normal
- Insisted as soon as he felt stable that he needed a step down – never actually stopped using opioids – would not “listen to reason”
- Relapsed, spent entire inheritance, back to OAT





# Involuntary Withdrawal

- Staff feel that for overall risk/benefit situation or safety reasons that patient should not continue with program
- Examples
  - - proven diversion of methadone or buprenorphine/naloxone
  - - threatens staff, pharmacist or other patients
  - - reckless use of benzodiazepines, opioids, alcohol
  - - erratic attendance at clinic

A dark blue arrow points to the right from the top left corner. Several thin, curved lines in shades of blue and grey sweep across the left side of the slide, starting from the bottom and curving upwards and to the right.

# Involuntary Withdrawal

- Clinics have different thresholds (depending on resource and staffing level) – staff should foster common principles and be able to discuss cases
- Offer the patient referral to a different OAT clinic, if possible and reasonable

A decorative graphic on the left side of the slide. It features a dark grey arrow pointing to the right at the top. Below the arrow, several thin, curved lines in shades of blue and grey sweep downwards and to the right, creating a dynamic, abstract background element.

## Pace of Involuntary Stepdown

- Rapid or slow - depends – may use outside pharmacy - warn pharmacy!!!
- If aggressive or difficult behaviour persists, rapidly stop treatment
- Actual threats or aggressive behaviour – consider police report

A decorative graphic on the left side of the slide. It features a dark grey arrow pointing right at the top, with several thin, curved lines in shades of blue and grey extending downwards from the arrow's tail.

## Jay

- Bright, always argumentative – fighting the rules – behaviour contract on chart
- Used foul threatening language to his case manager – team decision for involuntary withdrawal
- Rapid step down, at outside pharmacy
- Given name of alternate clinic and within 2 weeks enrolled there – “I know I need it”

A decorative graphic on the left side of the slide. It features a dark blue vertical bar on the far left. A black arrow points to the right from the top of this bar. Below the arrow, several thin, light blue lines curve downwards and to the right, creating a sense of movement and depth.

# Requests Return to Program?

- Clinic staff decide – behaviour is sometimes much better – but some clients are so difficult, return to program will not occur
- A waiting period may be beneficial – patient recognizes consequences are definitely in place



## Just Wait....

- ▶ They often return older and wiser and ready to work – and ready to follow the rules



**Questions?**